

Synthesis of Chiral Macrocycles: V.¹ Synthesis of Some *cyclo-(N^α-Dinicotinoyl)aromatic Octapeptides* and *cyclo-(N^α-Dinicotinoyl)pentapeptide Lysine Schiff Bases*²

A. M. Naglah^{a,b}, A. E. Amr^{a,c}, N. M. Khalifa^{a,d}, and M. A. Al-Omar^a

^a Pharmaceutical Chemistry Department, Drug Exploration & Development Chair (DEDC), College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia
e-mail: aeamr1963@yahoo.com

^b Peptide Chemistry Department, Chemical Industries Research Division, National Research Centre, 12622-Dokki, Cairo, Egypt

^c Applied Organic Chemistry Department, National Research Center, Cairo, Dokki 12622, Egypt

^d Therapeutic Chemistry Department, National Research Centre, 12622-Dokki, Cairo, Egypt

Received May 17, 2015

Abstract—A series of octa bridged peptide and macrocyclic pentapeptide Schiff base derivatives were synthesized from tetrapeptide hydrazide and macrocyclic pentapeptide ester. Condensation of hydrazide with acid anhydrides upon refluxing in acetic acid afforded the corresponding 3,5-bis-tetraimide pyridine and macrocyclic octapeptide derivatives. Treatment of macrocyclic pentapeptide ester with hydrazine hydrate gave the corresponding acid hydrazide derivative. Condensation of the latter with aromatic aldehydes led to the corresponding cyclic pentapeptide Schiff bases.

Keywords: synthesis, macrocyclic penta- and octa peptides, Schiff bases

DOI: 10.1134/S1070363215120324