

# CONTENTS

VOLUME 50

Number 3

2006

## RUSSIAN MATHEMATICS (Iz. VUZ)

|                                                                                                                                                                                                               | Pages<br>Russian/English |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------|
| <i>Weak presentations of computable ordered semigroups.</i>                                                                                                                                                   |                          |
| M. M. Arslanov and N. Kehayopulu . . . . .                                                                                                                                                                    | 3      1                 |
| <i>Operators of the multiplicative Fourier transform.</i> M.S. Bepalov . . . . .                                                                                                                              | 9      7                 |
| <i>Estimates of solutions of nonhomogeneous difference-differential equations of the neutral type.</i> V.V. Vlasov and S.A. Ivanov . . . . .                                                                  | 24      22               |
| <i>Extension of the definition domain of singular integral operators with degenerating symbol.</i> G.V. Kovaleva and <span style="border: 1px solid black; padding: 0 2px;">N.Ya. Tikhonenko</span> . . . . . | 31      29               |
| <i>On definability of derived objects on universal algebras.</i> A.G. Pinus . . . . .                                                                                                                         | 36      34               |
| <i>Two-sided estimates of <math>L^p</math>-norms of the stress function for convex domains in <math>\mathbb{R}^n</math>.</i>                                                                                  |                          |
| R.G. Salahudinov . . . . .                                                                                                                                                                                    | 41      39               |
| <i>Correct definition of initial-boundary-value problems for the generalized Kelvin–Voigt model.</i> M.V. Turbin . . . . .                                                                                    | 50      47               |
| <i>About a sufficient existence condition of a solution to the equation with nonmonotonic operators.</i> B.A. Shuvar and M.I. Kopach . . . . .                                                                | 59      56               |
| <i>On the Parsons model in the case of bifurcation point with two-dimensional branching.</i>                                                                                                                  |                          |
| L.D. Eskin . . . . .                                                                                                                                                                                          | 62      59               |
| <i>One hyperbolic equation in a three-dimensional space.</i> A.A. Kungurtsev . . . . .                                                                                                                        | 76      72               |